

## ACI OF FEMORAL CONDYLE REHABILITATION PROTOCOL

	<b>WEIGHT BEARING</b>	<b>BRACE</b>	<b>ROM</b>	<b>EXERCISES</b>
<b>PHASE I</b> 0-6 weeks	Non-WB	<p><b>0-2 weeks:</b> Locked in full extension at all times</p> <p>Off for CPM and exercise only</p> <p>Discontinue at 2 wks assuming quads can control SLR w/o extension lag</p>	<p><b>0-6 weeks:</b> Use CPM for 6 hours/day, beginning at 0-40°; advance 5-10° daily as tolerated</p>	<p><b>0-2 weeks:</b> Quad sets, SLR, calf pumps, passive leg hangs to 90° at home</p> <p><b>2-6 weeks:</b> PROM/AAROM to tolerance, patella and tibiofibular joint mobs, quad, hamstring, and glute sets, SLR, side-lying hip and core</p>
<b>PHASE II</b> 6-8 weeks	Advance 25% weekly until full	None	Full	Advance Phase I exercises
<b>PHASE III</b> 8-12 weeks	Full	None	Full	<p>Gait training, begin closed chain activities: wall sits, shuttle, mini-squats, toe raises</p> <p>Begin unilateral stance activities, balance training</p>
<b>PHASE IV</b> 12 weeks – 6 months	Full	None	Full	<p>Advance Phase III exercises; maximize core/glutes, pelvic stability work, eccentric hamstrings</p> <p>May advance to elliptical, bike, pool as tolerated</p>
<b>PHASE V</b> 6-12 months	Full	None	Full	<p>Advance functional activity</p> <p>Return to sport-specific activity and impact when cleared by MD after 8 mos</p>