## ACL PATELLAR TENDON AUTOGRAFT RECONSTRUCTION REHABILITATION PROTOCOL

	WEIGHT BEARING	BRACE	ROM	EXERCISES
PHASE I 0-4 weeks	Full in Brace	<ul> <li>0-1 week: Locked in full extension for ambulation and sleeping</li> <li>1-4 weeks: Unlocked for ambulation, remove for sleeping**</li> </ul>	As tolerated	Heel slides, quad/hamstring sets, patellar mobs, gastroc/soleus stretch SLR w/ brace in full extension until quad strength prevents extension lag Side-lying hip/core
PHASE II 4-12 weeks	Full	Discontinue at day 28 if patient has no extension lag	Full	Begin toe raises, closed chain quads, balance exercises, hamstring curls, stationary bike, step-ups, front and side planks; advance hip/core
PHASE III 12-16 weeks	Full	None	Full	Advance closed chain strengthening Progress proprioception activities Begin stairmaster, elliptical at 8 wks; Running straight ahead at 12 weeks
PHASE IV 16-24 weeks	Full	None	Full	<ul> <li>16 wks: Begin jumping</li> <li>20 wks: Advance to sprinting, backward running, cutting/pivoting/changing direction, initiate plyometric program and sport-specific drills</li> <li>22 wks: Advance as tolerated FSA completed at 22 wks***</li> </ul>
PHASE V > 6 months	Full	None	Full and pain- free	Gradual return to sports participation after completion of FSA Maintenance program based on FSA

\*Modified with concomitantly performed meniscus repair/transplantation or articular cartilage procedure

\*\*Brace may be removed for sleeping after first post-operative visit (day 7-10)

\*\*\*Completion of FSA (Functional Sports Assessment) not mandatory, but recommended at approx 22 wks post-op for competitive athletes returning to play after rehab